

Set Theory (Math 112)

First Midterm

Spring 2005

Ali Nesin

June 29, 2005

Let X be a set. A **filter** on X is a set \mathfrak{R} of subsets of X that satisfies the following properties:

- i) If $A \in \mathfrak{R}$ and $A \subseteq B \subseteq X$, then $B \in \mathfrak{R}$.
- ii) If A and B are in \mathfrak{R} , then so is $A \cap B$.
- iii) $\emptyset \notin \mathfrak{R}$ and $X \in \mathfrak{R}$.

If $A \subseteq X$ is a fixed nonempty subset of X , then the set of subsets $\mathfrak{R}(A)$ of X that contain A is a filter on X . Such a filter is called **principal filter**. If X is infinite, then the set of cofinite subsets of X is a filter, called **Fréchet filter**. A filter is called **ultrafilter** if it is a maximal filter.

We fix a set X .

1. Show that a principal filter $\mathfrak{R}(A)$ on X is an ultrafilter if and only if A is a singleton set.
2. Show that the Fréchet filter (on an infinite set) is not contained in a principal filter.
3. Show that the intersection of a set of filters is a filter.
4. Show that if \mathfrak{R} is a set of subsets of X such that $A_1 \cap A_2 \cap \dots \cap A_n \neq \emptyset$ for any $A_1, A_2, \dots, A_n \in \mathfrak{R}$, then there is a filter that contains \mathfrak{R} . Describe this filter in terms of \mathfrak{R} .
5. Show that a filter \mathfrak{R} is an ultrafilter if and only if for any $A \subseteq X$, either A or A^c is in \mathfrak{R} .
6. Conclude that any ultrafilter on X that contains a finite subset of X is a principal filter. Deduce that every nonprincipal ultrafilter contains the Fréchet filter.
7. (AC) Show that for any filter \mathfrak{R} on X , there is an ultrafilter on X that contains \mathfrak{R} .
8. (AC) Show that if X is infinite then there are nonprincipal ultrafilters on X .

9. Let \mathfrak{R} be a filter on X . Let A_x ($x \in X$) be sets. Consider the product

$$\begin{aligned} \prod_{x \in X} A_x &:= \{f = (f_x)_{x \in X} : f_x \in A_x \text{ for all } x \in X\} \\ &= \{f : X \longrightarrow \prod_{x \in X} A_x : f(x) \in A_x \text{ for all } x \in X\}. \end{aligned}$$

On the set $\prod_{x \in X} A_x$ consider the relation defined by

$$f \equiv g \iff \{x \in X : f(x) = g(x)\} \in \mathfrak{R}.$$

Show that this is an equivalence relation.

10. Show that if $A_x = A$ for all $x \in X$ and if \mathfrak{R} is nonprincipal filter then the map that sends an element $a \in A$ to the class of the constant function a is an injection from A into M .
11. Suppose that each A_x is a set ordered by a relation $<_x$. On M define the relation $<$ by,

$$[(f_x)_{x \in X}] < [(g_x)_{x \in X}] \iff \{x \in X : f_x <_x g_x\} \in \mathfrak{R}.$$

Show that this defines an order on M .

12. Show that if \mathfrak{R} is an ultrafilter and if each $(A_x, <_x)$ is totally ordered then so is $(M, <)$.
13. Suppose that each $(A_x, <_x)$ has a largest element. Is it true that $(M, <)$ has a largest element?
14. Suppose that each $(A_x, <_x)$ is well-ordered. Is it true that $(M, <)$ is well-ordered?