## Algebra Final Fall 2013 Ali Nesin

Throughout G denotes a group.

For  $X, Y \subseteq G$ , let [X, Y] denote the subgroup generated by all the elements of the form

$$[x, y] := x^{-1}y^{-1}xy$$

for  $x \in X$  and  $y \in Y$ .

- 1. Show that [X, Y] = [Y, X].
- 2. Show that if *H* and *K* are subgroups of *G* that normalize each other then  $[H, K] \leq H \cap K$ .
- 3. Show that for  $x, y, z \in G$ ,  $[x, yz] = [x, z][x, y]^z$  and  $[xy, z] = [x, z]^y[y, z]$ . Conclude that if  $H, K \leq G$ , then H and K normalize the subgroup [H, K]. Conclude also that if  $A \leq G$  is an abelian subgroup and if  $g \in N_G(A)$ , then the map  $ad(g) : A \to A$  defined by ad(g)(a) = [a, g] is a group homomorphism whose kernel is  $C_A(g)$ .

We define  $G^n$  and  $G^{(n)}$  by induction on n:  $G^0 = G^{(0)} = G$ ,  $G^{n+1} = [G, G^n]$ ,  $G^{(n+1)} = [G^{(n)}, G^{(n)}]$ . We let  $G' = G^1 = G^{(1)}$ .

- 4. Show that  $G^{n+1} \le G^n$  and that  $G^{(n+1)} \le G^{(n)}$ . Show also that  $G^n$  and  $G^{(n)}$  are characteristic subgroups of G.
- 5. Show that if  $H \triangleleft G$  and G/H is abelian then  $G' \subseteq H$ . Conversely show that if  $G' \subseteq H \subseteq G$ , then  $H \triangleleft G$  and G/H is abelian.
  - 6. Show that  $[G^i, G^j] \le G^{i+j+1}$  and  $G^{(i)} \le G^i$  for all i, j.

We define  $Z_n(G)$  by induction on n as follows:  $Z_n(G) = 1$  if  $n \le 0$  and for  $n \ge 0$ ,  $Z_{n+1}(G)$  is the unique subgroup of G that contains  $Z_n(G)$  such that  $Z(G/Z_n(G)) = Z_{n+1}(G)/Z_n(G)$ .

- 7. Show that  $Z_n(G)$  is a characteristic subgroup of G for all n.
- 8. Show that  $[G^i, Z_i] \le Z_{i-i-1}$  and that  $[Z^{i+1}, G^i] = 1$  for all i, j.

A group is said to be *solvable* if  $G^{(n)} = 1$  for some n. If  $G^{(n)} = 1$  but that  $G^{(n-1)} \neq 1$ , G is said to be solvable of class n.

A group is said to be *nilpotent* if  $G^n = 1$  for some n. If  $G^n = 1$  but that  $G^{n-1} \neq 1$ , G is said to be *nilpotent of class n*.

- 9. Show that a nilpotent group is solvable.
- 10. Let G be nilpotent of class n. Show that  $G^{n-i} \leq Z_i$ . Conclude that  $G = Z_n$ .
- 11. Conversely, assume that  $G = \mathbb{Z}_n$ . Show that  $G^i \leq \mathbb{Z}_{n-i}$ . Conclude that G is nilpotent of class n.
  - 12. Show that *G* is nilpotent of class *n* if and only if  $Z_n = G$  and  $Z_{n-1} \neq G$ .